

Section 31.9: cont'd:

Tuesday, April 23, 2013
11:35 AM

example: solve:

$$y'' - 5y' + 4y = 12e^x$$

complementary solution:

$$m^2 - 5m + 4 = 0$$
$$(m - 4)(m - 1) = 0$$
$$m = 4, 1$$

$$y_c = C_1 e^x + C_2 e^{4x}$$

particular solution:

~~$y_p = Ae^x$~~ ← bad case because Ae^x is a like term to one of the terms in y_c

$y_p = Axe^x$ ← so we fixed this by multiplying by x until no like terms

take derivatives, plug into DE:

$$y_p = Axe^x$$
$$y_p' = \downarrow \quad \searrow \quad Ae^x$$
$$y_p' = Axe^x + Ae^x$$
$$y_p'' = \downarrow \quad \searrow \quad Ae^x$$
$$y_p'' = Axe^x + Ae^x + Ae^x$$
$$= Axe^x + 2Ae^x$$

original DE:

original DE:

$$y'' - 5y' + 4y = 12e^x$$

$$\cancel{Ax^2e^x} + \cancel{2Ax}e^x - \cancel{5Ax}e^x - \cancel{5A}e^x + 4Ax^2e^x = 12e^x$$

$$-3Ae^x = 12e^x$$

$$-3A = 12$$

$$A = -4$$

$$y_p = -4xe^x$$

full solution:

$$y = y_c + y_p$$

$$= c_1 e^x + c_2 e^{4x} - 4xe^x$$

→ general solution